

Research Paper

Effect of cooking on nutritive and antioxidant properties of leafy vegetables consumed in Southern Côte d'Ivoire

Florentin C. Acho, *Lessoy T. Zoué, Gisèle Y. A. Koua, Sévérin A.K. Kra and Sébastien L. Niamké

Laboratoire de Biotechnologies, UFR Biosciences, Université Félix Houphouët-Boigny, 22 BP 582, Abidjan 22, Côte d'Ivoire

(Received April 15, 2014, Accepted July 05, 2014)

Abstract

African leafy vegetables are valuable sources of nutrients especially in rural areas where they contribute substantially to protein, minerals, vitamins, fibre and other beneficial nutrients. Five leafy vegetable species (*Basella alba*, *Colocasia esculenta*, *Corchorus olitorius*, *Solanum melongena* and *Talinum triangulare*) that are used as soup condiments in Southern Côte d'Ivoire were subjected to cooking in order to evaluate the effect of this processing method on their nutritive value and antioxidant properties. The result of the study revealed that longer time of cooking (higher than 15 min) caused negative impact by reducing nutritive value but positive impact by reducing anti-nutrients such as oxalates and phytates. The registered losses at 15 min were as follow: ash (12.42 – 54.09 %), proteins (3.47 - 33.23 %), vitamin C (46.42 – 83.21 %), carotenoids (73.74 – 100 %) oxalates (19.29 – 59.86 %) and phytates (28.61 – 78.02 %). The average reduction of polyphenols content at 15 min of cooking was 9.95 – 59.43 %. Contrary to these reductions, a slight increase of fibres content was observed in the studied cooked leafy vegetables. Furthermore after 15 min of cooking time the residual contents of minerals were: calcium (119.92 – 512.70 mg/100g), magnesium (51.94 – 285.07 mg/100g), potassium (529.92 – 1721.18 mg/100g), iron (12.37 – 25.63 mg/100g) and zinc (11.53 – 25.03 mg/100g). All these results suggest that the recommended time of domestic cooking must be less than 15 min for the studied leafy vegetables in order to contribute efficiently to the nutritional requirement and to the food security of Ivorian population.

Keywords: antioxidant properties, cooking processing, leafy vegetables, nutritive value.

Introduction

Leafy vegetables constitute essential components of human diet in Africa generally and particularly in West Africa ^[1,2,3,4]. These plants are important protective foods and highly beneficial for the maintenance of health and prevention of diseases as they contain valuable food ingredients which can be utilized to build up and repair the body ^[5]. Indeed, these plants are valuable sources of nutrients especially in rural areas where they contribute substantially to protein, minerals, vitamins, fibre and other nutrients which are usually in short supply in daily diets ^[6]. Antinutrients such as nitrates, phytates, tannins and oxalates have been reported in leafy vegetables ^[7,2,3,4]. Antinutritional factor is known to interfere with metabolic processes such that growth and bioavailability of nutrients are negatively influenced ^[8]. Phytates and oxalates have the ability to form chelates with di-valent metallic ions such as Ca, Mg, Zn and Fe to form poorly soluble compounds that are not readily absorbed from the gastrointestinal tract, thus decreasing their bioavailability ^[9]. In order to reduce these antinutritional factor contents, African leafy vegetables are usually blanched and cooked for soups preparation. Indeed, these vegetables are cooked and eaten as a relish together with a starchy

staple food, usually in the form of porridge ^[10]. Leafy vegetable (LV) dishes may be prepared from a single plant species or from a combination of different species, and the combinations eaten vary daily ^[11]. Ingredients such as salt, oil, butter, groundnuts, tomato and onion are usually added while cooking African leafy vegetables in order to enhance flavour, taste, color and aesthetic appeal ^[12]. These added ingredients also depend on availability and preference of consumers ^[13,14]. Several works and reports have revealed that cooking or preparation methods and period of cooking may affect the nutritional value as well as the bioavailability of many nutrients in leafy vegetables ^[15,16].

In Côte d'Ivoire, more than twenty (20) species of leafy vegetables belong to 6 botanical families, are consumed by populations through confectionary soups using boiling or blanched processing ^[17]. Furthermore, the consumption of these leafy vegetables is linked to the region and ethno-botanical studies have stated that most people in Southern Côte d'Ivoire consume indigenous green leafy vegetables such as *Basella alba* "epinard", *Colocasia esculenta* "taro", *Corchorus olitorius* "kplala", *Solanum melongena* "aubergine" and *Talinum triangulare* "mamichou" ^[18,19,20]. Earlier reports have highlighted the nutritive potential of these fresh leafy vegetables ^[2] but there is a lack of scientific data with regards to the effect of cooking methods on their physicochemical and nutritive characteristics. Therefore, the purpose of this study is to conduct investigation on the effect of cooking (boiling) on the nutritive value of these selected leafy vegetables in order to provide necessary informations for their wider utilization and contribution to food security of Ivorian populations.

Materials and Methods

Samples collection : Leafy vegetables (*Basella alba*, *Colocasia esculenta*, *Corchorus olitorius*, *Solanum melongena* and *Talinum triangulare*) were collected fresh and at maturity from cultivated farmlands located at Dabou (latitude: 5°19'14" North; longitude: 4°22'59"West) (Abidjan District). The samples were harvested at the early stage (between one and two weeks of the appearance of the leaves). These plants were previously authenticated by National Floristic Center (University Felix Houphouët-Boigny, Abidjan-Côte d'Ivoire).

Samples processing: The fresh leafy vegetables were destalked, washed with deionized water and allowed to drain at ambient temperature. Each sample was divided into two lots. The first lot (raw) was dried in an oven (Memmert, Germany) ^[21], ground with a laboratory crusher (Culatti, France) equipped with a 10 µm mesh sieve. Each sample was stored in a clean dry air-tight sample bottle in a refrigerator (4°C) until required for analyses. The second lot was cooked by using the method ^[22] and modified as follow: 250 g of leafy vegetables were immersed in 1.5 L of boiled water in stainless steel container for 15, 25 and 45 min. The cooking solution was discarded and the boiled samples were cooled, drained at ambient temperature and subjected to the same treatment using for raw samples.

Chemicals : All solvents (n-hexane, petroleum ether, acetone, ethanol and methanol) were purchased from Merck. Standards used (gallic acid, β-carotene) and reagents (metaphosphoric acid, vanillin, Folin-Ciocalteu, DPPH) were purchased from Sigma-Aldrich. All chemicals used in the study were of analytical grade.

Nutritive characterization

Physicochemical analysis: Ash, proteins and lipids were determined using official methods ^[23]. For crude fibres, 2 g of dried powdered sample were digested with 0.25 M sulphuric acid and 0.3 M sodium hydroxide solution. The insoluble residue obtained was washed with hot water and dried in an oven (Memmert, Germany) at 100 °C until constant weight. The dried residue was then incinerated, and weighed for the determination of crude fibres content. Carbohydrates and calorific value were calculated using the following formulas ^[24]:

Carbohydrates: $100 - (\% \text{ moisture} + \% \text{ proteins} + \% \text{ lipids} + \% \text{ ash} + \% \text{ fibres})$.

Calorific value: $(\% \text{ proteins} \times 2.44) + (\% \text{ carbohydrates} \times 3.57) + (\% \text{ lipids} \times 8.37)$.

The results of ash, fibres, proteins, lipids and carbohydrates contents were expressed on dry matter basis.

Oxalates and phytates determination: Oxalates content was performed using a titration method ^[25]. One (1) g of dried powdered sample was weighed into 100 mL conical flask. A quantity of 75 mL of

sulphuric acid (3 M) was added and stirred for 1 h with a magnetic stirrer. The mixture was filtered and 25 mL of the filtrate was titrated while hot against KMnO_4 solution (0.05 M) to the end point.

Phytates contents were determined using the Wade's reagent colorimetric method [26]. A quantity (1 g) of dried powdered sample was mixed with 20 mL of hydrochloric acid (0.65 N) and stirred for 12 h with a magnetic. The mixture was centrifuged at 12000 rpm for 40 min. An aliquot (0.5 mL) of supernatant was added with 3 mL of Wade's reagent. The reaction mixture was incubated for 15 min and absorbance was measured at 490 nm by using a spectrophotometer (PG Instruments, England). Phytates content was estimated using a calibration curve of sodium phytate (10 mg/mL) as standard.

Mineral analysis: The dried powdered samples (5 g) were burned to ashes in a muffle furnace (Pyrolabo, France). The ashes obtained were dissolved in 10 mL of HCl/ HNO_3 and transferred into 100 mL flasks and the volume was made up using deionized water. The mineral composition of each sample was determined using an Agilent 7500c inductively coupled argon plasma mass spectrometer (ICP-MS). Calibrations were performed using external standards prepared from a 1000 ppm single stock solution made up with 2% nitric acid

Antioxidant properties

Vitamin C and carotenoids determination: Vitamin C contained in analysed samples was determined by titration [27]. About 10 g of ground fresh leaves were soaked for 10 min in 40 mL metaphosphoric acid-acetic acid (2%, w/v). The mixture was centrifuged at 3000 rpm for 20 min and the supernatant obtained was diluted and adjusted with 50 mL of bi-distilled water. Ten (10) mL of this mixture was titrated to the end point with dichlorophenol-indophenol (DCPIP) 0.5 g/L.

Carotenoids were extracted and quantified by using a spectrophotometric method [28]. Two (2) g of ground fresh leaves were mixed three times with 50 mL of acetone until loss of pigmentation. The mixture obtained was filtered and total carotenoids were extracted with 100 mL of petroleum ether. Absorbance of extracted fraction was then read at 450 nm by using a spectrophotometer (PG Instruments, England). Total carotenoids content was subsequently estimated using a calibration curve of β -carotene (1 mg/mL) as standard.

Polyphenols determination: Polyphenols were extracted and determined using Folin–Ciocalteu's reagent [29]. A quantity (1 g) of dried powdered sample was soaked in 10 mL of methanol 70% (w/v) and centrifuged at 1000 rpm for 10 min. An aliquot (1 mL) of supernatant was oxidized with 1 mL of Folin–Ciocalteu's reagent and neutralized by 1 mL of 20% (w/v) sodium carbonate. The reaction mixture was incubated for 30 min at ambient temperature and absorbance was measured at 745 nm by using a spectrophotometer (PG Instruments, England). The polyphenols content was obtained using a calibration curve of gallic acid (1 mg/mL) as standard.

Antioxidant activity: Antioxidant activity assay was carried out using the 2,2-diphenyl-1-picrylhydrazyl (DPPH) spectrophotometric method [30]. About 1 mL of 0.3 mM DPPH solution in ethanol was added to 2.5 mL of sample solution (1 g of dried powdered sample mixed in 10 mL of methanol and filtered through Whatman No. 4 filter paper) and was allowed to react for 30 min at room temperature. Absorbance values were measured with a spectrophotometer (PG Instruments, England) set at 415 nm. The average absorbance values were converted to percentage antioxidant activity using the following formula:

$$\text{Antioxidant activity (\%)} = 100 - [(\text{Abs of sample} - \text{Abs of blank}) \times 100 / \text{Abs positive control}]$$

Statistical analysis: All the analyses were performed in triplicate and data were analyzed using EXCELL and STATISTICA 7.1 (StatSoft). Data were expressed as mean \pm standard deviation (SD).

Results and Discussion

Nutritive properties: The proximate composition of the cooked leafy vegetables examined in this study is presented in Table 1. The ash content after 15 min of cooking ranged from 6.90 ± 0.10 % (*C. esculenta*) to 14.50 ± 0.06 % (*T. triangulare*). These values were closed to 6.27 ± 0.12 % and 13.04 ± 0.12 % respectively, after 45 min of cooking for the same species. The decrease rate at 15 min (12.42 – 54.09 %) of ash content in the studied leafy vegetables may be a result of minerals leaching into the boiling water [31]. In spite of ash losses, the studied leafy vegetables may be considered as good

sources of minerals after 15 min of cooking when compared to values obtained for cereals and tubers^[32]. Cooking of all selected leafy vegetables resulted in a significant increase (1.37 – 21.67 %) in their crude fibres content at 15 min of cooking. This finding is in agreement with some reports that cooking caused increase in soluble fibres content, with corresponding decrease in insoluble fibres content^[33,34]. Indeed, cooking of plant tissues alters the physical and chemical properties of plant cell walls, which affects their performance as dietary fibre^[35]. The increased temperature during cooking leads to breakage of weak bonds between polysaccharides and the cleavage of glycosidic linkages, which may result in solubilization of the dietary fibre^[36].

With regard to their fibres content at 15 min (13.44 – 24.33 %), adequate intake of cooked leafy vegetables could lower the risk of hypertension, constipation, diabetes, colon and breast cancer^[37]. As concern proteins content, cooking processing used in this study caused 3.46 to 33.23 % reduction after 15 min. This reduced protein contents of cooked leafy vegetables could be attributed to the severity of thermal process during cooking to protein degradation^[38]. The relatively low level of proteins in cooked leafy vegetables necessitates supplementation with animal proteins or proteins from legumes to have an impact on protein-energy malnutrition^[39]. However, 15 min cooked leaves of *C. olitorius* could be considered as non negligible sources of proteins (17.79 ± 0.50 %) which content is higher than the minimal value (12%) recommended for protein foods^[40].

The relatively low values of lipids contents at 15 min of cooking (4.34 – 9.57 %) in the studied cooked leafy vegetables corroborate the findings of many authors which showed that leafy vegetables are poor sources of fat^[41]. The estimated calorific values (256.86 – 285.51 kcal/100 g) of the selected cooked leafy vegetables (15 min) compared favourably to 248.8 – 307.1 kcal/100 g reported in some Nigerian vegetables^[32]. Thus, the calorific values agree with general observation that vegetables have low energy values due to their low crude fat and relatively high level of moisture^[42]. This justify that cooked leafy vegetables are usually eaten as a relish together with a starchy staple food, usually in the form of porridge^[10].

The effect of cooking time on oxalates and phytates contents is depicted in figure 1. Levels losses at 15 min were 19.29 – 59.86 % and 26.81– 78.02 % for oxalates and phytates, respectively. At 45 min of cooking these values were 34.80 – 71.82 % and 36.67 – 85.30 %, respectively. The reductions in oxalates and phytates contents during cooking could be advantageous for improving the health status of consumers. Indeed, oxalates and phytates are anti-nutrients which chelate divalent cations such as calcium, magnesium, zinc and iron, thereby reducing their bioavailability^[43]. Therefore, cooking of leafy vegetables appears as a detoxification procedure by removing these anti-nutritional factors^[44].

Mineral composition of cooked leafy vegetables used in this study is shown in table 2. The residual contents of minerals after 15 min of cooking were: calcium (119.92 – 512.70 mg/100g), magnesium (51.94 – 285.07 mg/100g), potassium (529.92 – 1721.18 mg/100g), iron (12.37 – 25.63 mg/100g) and zinc (11.53 – 25.03 mg/100g). These observed reductions may be due to leaching of the mineral compounds into the boiling water^[45]. Considering the recommended dietary allowance (RDA) for minerals^[46]: calcium (1000 mg/day); magnesium (400 mg/day); iron (8 mg/day) and zinc (6 mg/day), consumption of 15 min cooked leafy vegetables could cover at least 50% RDA and could contribute for improving human diet. Calcium and phosphorus are associated for growth and maintenance of bones, teeth and muscles^[47]. As concern magnesium, this mineral is known to prevent cardiomyopathy, muscle degeneration, growth retardation, congenital malformations and bleeding disorders^[48].

Iron plays important role in prevention of anemia which affects more than one billion people worldwide^[49]. To predict the bioavailability of calcium and iron, anti-nutrients to nutrients ratios of cooked leafy vegetables were calculated (Table 3). The calculated [phytates]/[Ca] ratio in all the species were below the critical level of 2.5 whatever the cooking time. This critical value is known to impair calcium bioavailability^[50]. It was also observed that the calculated [phytates]/[Fe] ration for 15 min cooking time was above the critical level of 0.4 which is known to impair iron bioavailability^[51].

Table 1
Proximate composition of raw and cooked leafy vegetables consumed in Southern Côte d'Ivoire

	Ash (%)	Fibres (%)	Proteins (%)	Lipids (%)	Carbohydrates (%)	Calorific energy (kcal/100g)
<i>C. esculenta</i>						
Raw	15.03± 0.23	24.00± 0.46	9.80± 0.16	8.35± 0.15	42.85± 2.68	252.34± 14.55
15 min	6.90 ± 0.10	24.33± 0.52	9.46± 0.08	9.57± 0.13	49.40± 0.51	279.54± 1.24
25 min	6.60 ± 0.20	24.50± 1.30	9.21± 0.51	10.40± 0.26	49.29± 2.25	285.48± 4.59
45 min	6.27 ± 0.12	24.67± 0.38	8.33± 0.50	10.58± 0.03	50.15± 0.59	287.91± 1.44
<i>B. alba</i>						
Raw	19.79± 0.44	16.50± 0.30	9.86± 0.10	6.85± 0.05	47.00± 0.89	249.18± 4.41
15 min	12.47± 0.23	17.92± 0.72	8.87± 0.00	7.71± 0.29	53.03± 0.53	275.49± 4.07
25 min	12.1 ± 0.17	18.67± 0.29	6.83± 0.50	8.35± 0.31	50.82± 0.96	268.23± 4.96
45 min	10.73± 0.12	19.58± 0.72	6.33± 0.07	10.30± 0.72	53.06± 0.63	291.08± 6.06
<i>S. melongena</i>						
Raw	20.32± 2.36	13.70± 0.65	12.34± 0.09	2.73± 0.06	50.91± 3.16	277.57± 13.54
15 min	13.70 ±0.00	16.67± 0.93	11.37± 0.00	4.47± 0.21	53.70± 1.16	256.86± 3.78
25 min	13.19 ±0.00	20.00± 0.17	10.75± 0.00	5.24± 0.25	50.82± 0.31	251.51± 1.74
45 min	13.00 ±0.00	20.11± 0.42	7.58± 0.50	6.22± 0.05	53.06 ± 0.63	260.08± 1.07
<i>T. triangulare</i>						
Raw	22.20± 0.37	13.9 ± 1.50	17.18± 0.05	4.90± 0.06	52.77± 2.08	271.32± 8.17
15 min	14.50± 0.06	15.10 ±0.29	11.47± 0.17	5.95± 0.09	52.84± 0.59	266.42± 2.46
25 min	13.31± 0.10	16.17± 0.17	9.91± 0.51	7.77± 0.87	52.83± 1.02	277.81± 4.55
45 min	13.04± 0.12	16.28± 0.48	9.75± 0.00	8.70± 0.34	52.23± 0.74	283.07± 0.20
<i>C. olitorius</i>						
Raw	8.53 ± 0.15	11.49± 0.03	21.12± 0.05	3.28± 0.30	55.58± 0.84	277.40± 6.26
15 min	7.47 ± 0.12	13.44±0.20	17.79± 0.50	4.34± 0.30	56.96± 0.62	285.51± 1.24
25 min	7.27 ± 0.12	14.50± 1.32	15.46± 0.51	4.96± 0.36	57.81± 0.93	281.67± 4.77
45 min	6.87 ± 0.12	16.72± 0.39	13.41± 0.51	5.01± 0.20	57.99± 0.31	281.43± 2.61

Data are represented as means ± SD (n=3).

Table 2
Mineral composition (mg/100g) of raw and cooked leafy vegetables consumed in Southern Côte d'Ivoire

	Ca	Mg	P	K	Fe	Na	Zn
<i>C. esculenta</i>							
Raw	587.24 ± 0.55	347.29 ± 0.32	788.00 ± 0.74	2281.63 ± 2.14	143.37±0.13	39.45 ± 0.16	37.29 ± 0.03
15 min	119.92 ± 4.50	64.37±1.35	70.25 ± 3.80	529.92 ± 2.70	14.15 ± 0.90	12.20 ± 0.15	12.35 ± 0.10
25 min	107.42 ± 8.28	57.72 ± 1.56	67.20 ± 5.18	453.42 ± 3.96	9.10 ± 0.58	7.75 ± 0.30	10.23 ± 0.05
45 min	76.07 ± 3.35	30.23 ± 6.27	48.62 ± 2.54	329.17 ± 3.13	7.80 ± 0.14	5.09 ± 0.00	7.35 ± 0.03
<i>B. alba</i>							
Raw	750.34 ± 0.53	753.88 ± 0.53	390.14 ± 0.27	2709.43 ± 1.93	77.47 ± 0.05	555.01 ± 5.99	67.25 ± 0.72
15 min	375.71 ± 9.03	223.38 ± 8.71	95.15 ± 2.42	906.57 ± 6.14	12.37 ± 0.49	207.15 ± 0.60	25.03 ± 0.07
25 min	270.47 ± 2.21	195.36 ± 4.56	83.53 ± 2.92	794.97 ± 9.68	9.70 ± 0.05	127.35 ± 2.00	22.50 ± 0.67
45 min	162.81 ± 9.09	168.76 ± 1.39	54.93 ± 2.73	653.45 ± 54.21	7.90 ± 0.05	95.25 ± 0.25	17.05 ± 0.02
<i>S. melongena</i>							
Raw	796.54 ± 0.55	481.90 ± 0.33	374.46 ± 0.26	2256.10 ± 1.57	139.48±0.09	323.13 ± 2.96	64.64 ± 0.04
15 min	369.63 ± 3.75	223.75 ± 8.50	98.63 ± 1.00	885.50 ± 8.50	25.63 ± 1.38	165.00 ± 5.50	14.23 ± 0.01
25 min	327.25 ± 6.25	90.50 ± 6.50	87.50 ± 0.50	576.25 ± 8.50	15.00 ± 0.50	112.50 ± 2.50	12.62 ± 0.03
45 min	205.00± 1.25	70.62 ± 4.25	59.25 ± 5.25	393.75 ± 5.62	9.30 ± 0.25	95.62 ± 1.12	10.05 ± 0.02
<i>T. triangulare</i>							
Raw	601.37 ± 0.38	755.97 ± 0.48	239.59 ± 0.81	5053.23 ± 3.21	102.28±0.06	260.25 ± 0.75	36.10 ± 0.02
15 min	572.10 ± 9.14	285.07 ± 8.68	148.61 ± 4.57	1721.18 ± 10.73	13.09 ± 0.30	96.35 ± 0.80	15.20 ± 0.09
25 min	260.16 ± 9.93	224.52 ± 4.59	119.79 ± 6.65	1185.08 ± 48.58	9.86 ± 0.02	78.60 ± 0.70	13.65 ± 0.07
45 min	96.28 ± 6.08	134.48 ± 5.21	97.8 ± 6.52	1106.13 ± 10.40	6.29 ± 0.05	32.50 ± 0.00	11.90 ± 0.09
<i>C. olitorius</i>							
Raw	369.02 ± 2.33	234.51 ± 0.38	316.82 ± 0.52	2622.57 ± 16.56	97.60 ± 0.16	27.75 ± 0.08	24.71 ± 0.04
15 min	133.20 ± 6.48	51.94 ± 3.11	158.66 ± 7.35	1647.88 ± 47.06	19.42 ± 0.70	8.00 ± 0.00	11.53 ± 0.00
25 min	98.557± 2.52	39.30 ± 0.90	127.47 ± 3.81	1241.72 ± 5.08	10.90 ± 0.80	6.10 ± 0.00	9.77 ± 0.04
45 min	60.125 ± 5.87	29.78 ± 2.59	85.10 ± 3.35	1045.61 ± 30.61	6.18 ± 0.60	3.25 ± 0.00	7.05 ± 0.02

Data are represented as means ± SD (n=3).

Table 3: Anti-nutritional factors/mineral ratios of raw and cooked leafy vegetables consumed in Southern Côte d'Ivoire

	Phytates/Ca	Phytates/Fe	Oxalates/Ca
<i>C. esculenta</i>			
Raw	0.04	0.18	0.98
15 min	0.08	0.75	2.63
25 min	0.09	1.06	2.43
45 min	0.11	1.15	3.15
<i>B. alba</i>			
Raw	0.02	0.25	0.87
15 min	0.04	1.14	1.25
25 min	0.05	1.47	1.59
45 min	0.06	1.37	1.87
<i>S. melongena</i>			
Raw	0.05	0.29	0.12
15 min	0.04	0.70	0.10
25 min	0.05	1.19	0.11
45 min	0.08	1.75	0.13
<i>T. triangulare</i>			
Raw	0.05	0.28	0.86
15 min	0.01	0.49	0.73
25 min	0.02	0.50	1.51
45 min	0.04	0.68	3.52
<i>C. olerius</i>			
Raw	0.10	0.40	2.11
15 min	0.19	1.31	4.29
25 min	0.25	2.30	4.98
45 min	0.41	3.97	6.95

Antioxidant properties

Cooking resulted in a higher decrease of carotenoids and vitamin C contents in the studied leafy vegetables (Figure 2). For carotenoids, losses at 15 min were estimated to 73.74 to 100% when considering the cooking times. The decrease in the concentration of the total carotenoids observed during cooking could be attributed to wilting and damage of the leaf tissues which resulted in leaching. Also, excessive activity of lipoxygenase enzyme in the initial (warming) stages of cooking, oxidation and isomerization of β -carotene might account for the decrease in their concentration [52]. Carotenoids are considered as sources of provitamin A in plants and their amount determine their bioavailability in human diet [28].

Therefore, increased intake of cooked leafy with fat added, contributed significantly to improving the vitamin A status in children [53]. For vitamin C content, a significant reduction (46.42 – 83.21 %) was highlighted at 15 min during cooking processing (Figure 2). These values support the results obtained for other studies which indicate losses up to 66 % in cooked vegetables [54]. It is important noting that ascorbic acid is a water-soluble antioxidant that promotes absorption of soluble iron by chelating or by maintaining the iron in the reduced form [55]. With regard to the drastic decrease of vitamin C during cooking, consumption of cooked leafy vegetables may be supplemented with other sources of vitamin C such as tropical fruits to cover the daily need for humans (40 mg/day) as recommended by food agriculture organization [46].

The effect of cooking on polyphenols content and antioxidant activity of the selected leafy vegetables is depicted in figure 3. The losses of polyphenols contents at 15 min of cooking were 9.95 – 59.43 %. The decrease of the polyphenolic compounds content and overall antioxidant activity could be due to leaching or heat lability of specific flavonoids [56]. Indeed, flavonoids such as myricetin, quercetin, kaempferol, isorhamnetin and luteolin have been previously reported in leafy vegetables [57]. The negative impact of cooking on polyphenols content may affect the medicinal potentialities of leafy vegetables as previously mentioned [5]. Therefore, other processing methods such as blanching, drying, roasting and microwaving of leafy vegetables should be used to limit or avoid losses of polyphenols by leaching as observed during cooking.






Figure 1: Effect of cooking time on oxalates (A) and phytates (B) contents of leafy vegetables consumed in Southern Côte d'Ivoire






Figure 2: Effect of cooking time on vitamin C (A) and carotenoids (B) contents of leafy vegetables consumed in Southern Côte d'Ivoire





Figure 3: Effect of cooking time on polyphenols content (A) and antioxidant activity (B) of leafy vegetables consumed in Southern Côte d'Ivoire

Conclusion

African leafy vegetables (ALVs) contain significant levels of nutrients that are essential for human health. The result of this study revealed that leafy vegetables consumed in Southern Côte d'Ivoire contain appreciable amount of minerals, polyphenols and antinutrients. Cooking at 15, 25 and 45 minutes decreased considerably the nutritional value of these leafy vegetables. Nevertheless, the losses in antinutrients (oxalates, phytates) might have asserted a beneficial effect on bioavailability of minerals like iron and calcium. Thus, the study suggests that the recommended time of domestic cooking must be less than 15 min for the studied leafy vegetables in order to contribute efficiently to the nutritional requirement and to the food security of Ivorian population.

References

1. Kubmarawa D., Andenyang I. F. and Magomya A. M.; Proximate composition and amino acid profile of two non-conventional leafy vegetables (*Hibiscus cannabinus* and *Haematostaphis barteri*) Afr. J. Food Sci. 3, 233-236, (2009).
- 2 Constant F. Acho, Lessoy T. Zoué, Eric E. Akpa, Valery G. Yapo and Sébastien L. Niamké, Leafy vegetables consumed in Southern Côte d'Ivoire: a source of high value nutrients. J. Anim. Plant Sci. 20, 3159-3170, (2014).
- 3 Oulai Patricia, Lessoy Zoué, Rose-Monde Mégnanou, Ryta Doué and Sébastien Niamké, Proximate composition and nutritive value of leafy vegetables consumed in Northern Côte d'Ivoire. Eur. Sci. J., 10, 212-227, (2014).
4. Armel F. Zoro, Lessoy T. Zoué, Sévérin A.K. Kra, Arnaud E. Yépié and Sébastien L. Niamké, An overview of nutritive potential of leafy vegetables consumed in Western Côte d'Ivoire. Pak. J. Nutr. 12, 949-956, (2013).
5. Falade O. S., Sowunmi O. R., Oladipo A., Tubosun A. and Adewusi S.R.A.; The level of organic acids in some Nigerian fruits and their effects on mineral availability in composite diets. Pak. J. Nutr., 2, 82-88, (2003).
6. Mohammed M. I. and Sharif N.; Mineral composition of some leafy vegetables consumed in Kano, Nigeria. Nig. J. Basic Appl. Sci. 19, 208-211, (2011).
7. Besche T., Fretzdorff B. and Langenkemper G., Health protection by decreasing oxalate and increasing ascorbate in vegetables using biotechnology or gene transfer, sub-project. Alliance project and nutrition research institute for biochemistry of cereals and potatoes at the federal centre for cereal potato and lipid research. Detmold, Germany, (1988).
8. Akindahunsi A. A. and Salawu S. O., Phytochemical screening and nutrient and anti-nutrient composition of selected tropical green leafy vegetables. Afr. J. Biotechnol. 4, 497-501, (2005).
9. Soetan K.O. and Oyewole O. E., The need for adequate processing to reduce the anti-nutritional factors in plants used as human foods and animal feeds: A review. Afr. J. Food Sci., 9, 223-232, (2009).
10. Vainio-Mattila K.; Wild vegetables used by the Sambia in the Usambara Mountains, North-East Tanzania. Annales Botanici Fennici. 37, 57-67, (2000).
11. Marshall F.; Agriculture and use of wild and weedy greens by the Piik Ap Oom Okiek of Kenya. Econ. Botany. 55, 32-46, (2001).
12. Mepba H. D., Eboh L. and Banigo D. E. B., Effects of processing treatments on the nutritive composition and consumer acceptance of some Nigerian edible leafy vegetables. AJFAND. 1, 1-18, (2007).
13. Nguni D. and Mwila G., Opportunities for increased production, utilization and income generation from African leafy vegetables in Zambia. Retrieved September 2008 from the African Journal of Food, Agriculture, Nutrition and Development on the World Wide Web: <http://www.ajfand.net/>, (2007).

14. Ogoye-Ndegwa C. and Aagaard-Hansen J., Traditional gathering of wild vegetables among the Luo of Western Kenya – a nutritional anthropology project. *Ecol. Food Nutr.* 42, 69–89, **(2003)**.
15. Gupta S. and Bains K., Traditional cooked vegetable dishes as important sources of ascorbic acid and β -carotene in the diets of Indian urban and rural families. *Food Nutr. Bull.* 27, 306–310, **(2006)**.
16. Leskova E., Kobikova J., Kovacikova E., Kosicka M., Porubska J. and Holcikova K., Vitamin losses: retention during heat treatment and continual changes expressed by mathematical models. *J. Food Comp. Anal.* 19, 252–276, **(2006)**.
17. CNRA. Socio-economical importance of leafy vegetables for the urban populations of Côte d'Ivoire, CNRA Ed., **(2011)**.
18. Kouamé N. M., Contribution à l'étude des plantes spontanées alimentaires du département d'Oumé (Côte d'Ivoire). Mémoire de D.E.A d'Ecologie tropicale. Université de Cocody-Abidjan, Côte d'Ivoire, Pp. 122 **(2000)**.
19. Fondio L., Kouamé C., N'zi J. C., Mahyao A., Agbo E. and Djidji A. H., Survey of indigenous leafy vegetables in the urban and peri-urban areas of Côte d'Ivoire. In: M. L. Chadha et al. (Eds). *Indigenous vegetables and legumes: prospects for fighting poverty, hunger and malnutrition. Acta Horticulturae.* 752, 287-289, **(2007)**.
20. N'Dri M. T., Kouamé G. M., Konan E. and Traoré D., Plantes alimentaires spontanées de la région du Fromager (Centre-Ouest de la Côte d'Ivoire) : flore, habitats et organes consommés. *Sci. Nat.* 1:, 61-70, **(2008)**.
21. Chinma C. E. and Igyor M. A., Micro-nutriments and anti-nutritional content of select tropical vegetables grown in south-east, Nigeria. *Nig. Food J.* 25, 111-115, **(2007)**.
22. Randrianatoandro V. A., Identification et caractérisation des plats sources en micronutriments consommés en milieu urbain (Manjakaray, Madagascar) : étude des plats à base de légumes-feuilles, Thèse, Université d'Antananarivo, Madagascar, Pp. 150 **(2010)**.
23. AOAC., Official methods of analysis. Association of Official Analytical Chemists Ed., Washington DC, 684 **(1990)**.
24. FAO., Food energy-methods of analysis and conversion factors. FAO Ed, Rome, 97 **(2002)**.
25. Day R. A. and Underwood A. L., Quantitative analysis. 5th ed. Prentice Hall. 701 **(1986)**.
26. Latta M. and Eskin M., A simple method for phytate determination. *J. Agric. Food Chem.* 28, 1313-1315, **(1980)**.
27. Pongracz G., Weiser H. and Matzinger D., Tocopherols- Antioxydant. *Fat Sci. Technol.* 97, 90-104, **(1971)**.
28. Rodriguez-Amaya D. B., A guide to carotenoids analysis in foods. ILSI Press, Washington DC, 64 **(2001)**.
29. Singleton V. L., Orthofer R. and Lamuela-Raventos R. M., Analysis of total phenols and other oxydant substrates and antioxydants by means of Folin-ciocalteu reagent. *Methods Enzymol.* 299, 152-178, **(1999)**.
30. Choi C. W., Kim S. C., Hwang S. S., Choi B. K., Ahn H. J., Lee M. Z, Park S. H. and Kim S. K., Antioxydant activity and free radical scavenging capacity between Korean medicinal plant and flavonoids by assay guided comparison. *Plant Sci.* 163, 1161-1168, **(2002)**.
- 31 Oboh G., Effect of blanching on the antioxidant properties of some tropical green leafy vegetables. *Food Sci. Tec.* 38, 513-517, **(2005)**.

32. Antia B. S., Akpan E. J., Okon P. A. and Umoren I. U., Nutritive and antinutritive evaluation of sweet potato (*Ipomea batatas*) leaves. Pak. J. Nutr. 5, 166-168, **(2006)**.
33. Slavin J. L., Dietary fiber and body weight. J. Am. Diet Ass. 87, 1164-1168, **(1987)**.
34. Lintas C. and Cappelloni M., Dietary fiber, resistant starch and in vitro starch digestibility of cereal meals. Food Sci. Nutr. 42, 117 – 124, **(1998)**.
35. McDougall G. J., Morrison I. M., Stewart D. and Hillman J. R., Plant cell walls as dietary fibre: range, structure, processing and function. J. Sci. Food Agric. 70, 133–150, **(1996)**.
36. Svanberg S. M., Nyman E. M. G., Andersson L. and Nilsson R., Effects of boiling and storage on dietary fiber and digestible carbohydrates in various cultivars of carrots. J. Sci. Food Agric. 73, 245–254, **(1997)**.
37. Ishida H., Suzuno H., Sugiyama N., Innami S., Todokoro T. and Maekawa A., 2000. Nutritional evaluation of chemical component of leaves stalks and stems of sweet potatoes (*Ipomea batatas*). Food Chem. 68, 359-367, **(2000)**.
38. Lund D. B., 1997. Effects of heat processing on nutrients. In: Nutritional Evaluation of Food Processing, (R. Harries and E. Karmas, eds). The AVI Publishing Co. Inc Westport, pp. 205 – 203, **(1997)**.
39. Nangula P. U., Andre O., Kwaku G. D., Megan J. B., Mieke F., Nutritional value of leafy vegetables of sub-Saharan Africa and their potential contribution to human health: A review. J. Food Comp. Anal. 23, 499–509, **(2010)**.
40. Ali A., Proximate and mineral composition of the marchubeh (*Asparagus officinalis*). World Dairy Food Sci. 4, 142-149, **(2009)**.
41. Ejoh A. R., Tchouanguap M. F. and Fokou E., 1996. Nutrient composition of the leaves and flowers of *Colocasia esculenta* and the fruits of *Solanum melongena*. Plant Food Hum. Nutr. 49, 107-112, **(1996)**.
42. Sobowale S. S., Olatidoye O. P., Olorode O. O. and Akinlotan J. V., Nutritional potentials and chemical value of some tropical leafy vegetables consumed in south west Nigeria J. Sci. Multidisciplinary Res. 3, 55-65, **(2011)**.
43. Sandberg A. S., Bioavailability of minerals in legumes. Brit. J. Nutr. 88, 281-285, **(2002)**.
44. Ekop A. S. and Eddy N. O., 2005. Comparative Studies of the level of toxicants in the seed of Indian almond (*Terminalia catappa*) and African walnut (*Coula edulis*). Chem. Class J. 2, 74-76, **(2005)**.
45. Ajala L., The effect of boiling on the nutrients and anti-nutrients in two non conventional vegetables. Pak. J. Nutr. 8, 1430-1433, **(2009)**.
46. FAO., Human vitamin and mineral requirements. FAO Ed, 361 **(2004)**.
47. Turan M., Kordali S., Zengin H., Dursun A. and Sezen Y., Macro and micro-mineral content of some wild edible leaves consumed in Eastern Anatolia. Plant Soil Sci, 53, 129-137, **(2003)**.
48. Chaturvedi V. C., Shrivastava R. and Upreti R. K., Viral infections and trace elements: A complex trace element. Curr. Sci. 87, 1536-1554, **(2004)**.
49. Trowbridge F. and Martorell M., Forging effective strategies to combat iron deficiency. J. Nutri. 85, 875-880, **(2002)**.
50. Hassan L. G., Umar K. J. and Umar Z., Antinutritive factors in *Tribulus terrestris* (Linn) leaves and predicted calcium and zinc bioavailability. J. Trop. Biosci. 7, 33-36, **(2007)**.

51. Umar K. J., Hassan L. G., Dangoggo S. M., Inuwa M. and. Amustapha M. N., Nutritional content of *Melochia corchorifolia* (Linn.) leaves. *Int. J. Biol. Chem.* 1, 250-255, **(2007)**.
52. Speek A. J., Temalilwa G. R., Schrijver J., Determination of β -carotene content and vitamin A activity of vegetables by HPLC. *Food Chem.* 19, 65-74, **(1986)**.
53. Takyi E.E.K., Children's consumption of dark green leafy vegetables with added fat enhances serum retinol. *J. Nutr.* 129, 1549–1554, **(1999)**.
54. Fafunso M. and Bassir O., Effect of cooking on vitamin C content of fresh leaves and wilted leaves. *J. Agric. Food Chem.* 23, 9-17, **(1987)**.
55. Yamaguchi T., Mizobuchi T., Kajinawa H., Miyabe F., Terao J., Takamura H. and Matoba T., Radical-scavenging activity of vegetables and the effect of cooking on their activity. *Food Sci. Technol. Res.* 7, 250–257, **(2001)**.
56. Trichopoulou A., Vasilopoulou E., Hollman P., Chamalides C., Foufa E., Kaloudis T., Kromhout D., Miskaki P., Petrochilou I., Poulima E., Stafilakis K. and Theophilou D., Nutritional composition and flavonoid content of edible wild greens and green pies: a potential rich source of antioxidant nutrients in the Mediterranean diet. *Food Chem.* 70, 319-323, **(2000)**.
57. Wong S. P., Leong L. P. and Koh J. H., Antioxidant activities of aqueous extracts of selected plants. *Food Chem.* 99, 775-783, **(2006)**.